

Table 2

Dietary fibre content of various food sources

Source	Dietary fibre (g/100 g edible portion)		
	Total	Insoluble	Soluble
Grains			
Barley	17.3	—	—
Corn	13.4	—	—
Oats	10.3	6.5	3.8
Rice (dry)	1.3	1.0	0.3
Rice (cooked)	0.7	0.7	0.0
Wheat (whole grain)	12.6	10.2	2.3
Wheat germ	14.0	12.9	1.1
Legumes & pulses			
Green beans	1.90	1.40	0.50
Soy	15.0	—	—
Peas, green frozen	3.5	3.2	0.3
Kidney beans, canned	6.3	4.7	1.6
Lentils, raw	11.4	10.3	1.1
Lima beans, canned	4.2	3.8	0.4
White beans, raw	17.7	13.4	4.3
Vegetables			
Potato, no skin	1.30	1.0	0.30
Bitter melon	16.6	13.5	3.1
Beetroot	7.8	5.4	2.4
Fenugreek leaves	4.9	4.2	0.7
Ladyfinger	4.3	3.0	1.3
Spinach, raw	2.6	2.1	0.5
Turnips	2.0	1.5	0.5
Tomato, raw	1.2	0.8	0.4
Green onions, raw	2.2	2.2	0.0
Eggplant	6.6	5.3	1.3
Cucumbers, peeled	0.6	0.5	0.1
Cauliflower, raw	1.8	1.1	0.7
Celery, raw	1.5	1.0	0.5

Source Farhath Khanum et al. [2000](#); Schakel et al. [2001](#)